

The Investigation and Treatment of Diabetic Gastroparesis

Mohit Kumar, MRCP¹; Adam Chapman, BMBS²; Saad Javed, MBChB³; Uazman Alam, MRCP, PhD^{4,5,6}; Rayaz A Malik, FRCP, PhD^{3,7}; and Shazli Azmi, MRCP, PhD^{3,8}

¹Royal Albert Edward Infirmary Wroughtington, Wigan and Leigh National Health Service Foundation Trust, Wigan, United Kingdom; ²Aintree University Hospital National Health Service Foundation Trust, Liverpool, United Kingdom; ³Institute of Cardiovascular Sciences, University of Manchester, Manchester, United Kingdom; ⁴Division of Endocrinology, Diabetes and Gastroenterology, University of Manchester, Manchester, United Kingdom; ⁵Diabetes and Endocrinology Research and the Pain Research Institute, Department of Eye and Vision Sciences, Institute of Ageing and Chronic Disease, University of Liverpool and Aintree University Hospital NHS Foundation Trust, Liverpool, United Kingdom; ⁶Department of Diabetes and Endocrinology, Royal Liverpool and Broadgreen University NHS Hospital Trust, Liverpool, United Kingdom; ⁷Weill Cornell Medicine-Qatar, Doha, Qatar; and ⁸Manchester Diabetes Centre, Manchester University Foundation Trust, Manchester, United Kingdom

ABSTRACT

Purpose: This review provides an update on the investigations and treatment options for gastroparesis.

Methods: A comprehensive literature search of Medline, PubMed, Embase and OVID was conducted which included all systematic reviews and research articles that focused on the diagnosis, investigations and management diabetic gastroparesis.

Findings: Dietary modifications and pharmacologic treatment with prokinetics to increase gastric motility form the mainstay of treatment. However, the use of prokinetics is limited by adverse effects and serious adverse effects, leaving metoclopramide as the only drug approved by the US Food and Drug Administration for the treatment of gastroparesis. Newer therapies, including motilin receptor agonists, ghrelin receptor agonists, and neurokinin receptor antagonists, are currently being investigated. Transpyloric stenting, gastric electrical stimulation, and gastric per-oral endoscopic myotomy provide mechanical options for intervention, and surgical interventions in severe intractable gastroparesis include laparoscopic pyloroplasty or gastrectomy.

Implications: Advances to better understand the pathophysiology and management of diabetic gastroparesis have been limited, especially with discordance between symptoms and severity of delay in gastric emptying. Established treatment options are limited;

however, recent pharmacologic and surgical interventions show promise. (*Clin Ther.* 2018;■:■■■-■■■) © 2018 Elsevier HS Journals, Inc. All rights reserved.

Key words: Type 1 Diabetes Mellitus, Type 2 Diabetes Mellitus, Gastroparesis, Microvascular.

INTRODUCTION

First described by Rundles in 1945,¹ the term *gastroparesis diabeticorum* was coined by Kassander in 1958.² The most common cause of gastroparesis, a condition characterized by delayed gastric emptying in the absence of a mechanical obstruction, is diabetes mellitus.^{3,4} This condition can manifest in a variety of symptoms, including early satiety, nausea, vomiting, and anorexia. It can be associated with significant morbidity and impaired quality of life, with anxiety and depression and an effect on patients' self-management of diabetes, especially with fluctuating glucose levels.^{5,6} However, the association between the severity of symptoms and delay in gastric emptying is not linear.⁷ Indeed, up to 40% of patients with

Accepted for publication April 17, 2018.

<https://doi.org/10.1016/j.clinthera.2018.04.012>

0149-2918/\$ - see front matter

© 2018 Elsevier HS Journals, Inc. All rights reserved.

diabetic gastroparesis (DG) can be asymptomatic.⁸ The term *gastric hypoglycemia* has been used to describe patients with hypoglycemia attributable to gastroparesis, and it should be considered in the differential diagnosis for patients diagnosed as having brittle diabetes.^{9,10} A comprehensive literature search of Medline, PubMed, Embase and OVID was conducted which included all systematic reviews and research articles that focused on the diagnosis, investigations and management of diabetic gastroparesis.

EPIDEMIOLOGY

The exact prevalence of DG remains unknown. A major population-based study, evaluating symptoms, reported a 10-year cumulative incidence of gastroparesis of 5.2% in patients with type 1 diabetes mellitus (T1DM), 1.0% in patients with type 2 diabetes mellitus (T2DM), and 0.2% in controls.¹¹ The T1DM Exchange registry data reported 4.8% of patients had gastroparesis, with significant associations with female sex, longer duration of diabetes, older age, and more frequent severe hypoglycemic episodes, despite higher glycosylated hemoglobin (HbA_{1c}) levels.¹² Although less common than in patients with T1DM, the greater prevalence of T2DM means that more patients have gastroparesis attributable to this condition.⁸ There are conflicting data on the prevalence of symptoms of gastroparesis in patients with diabetes.^{13,14} In addition to delayed gastric emptying, patients may also develop rapid gastric emptying, with one study reporting this phenomenon in 22% of patients evaluated with scintigraphy.¹⁵

An interesting phenomenon is that of symptom turnover (the appearance and disappearance of symptoms over time), and several studies found this to be common in patients with DG. One study remarked that the turnover of symptoms was associated with depression rather than other factors, such as glycemic control.^{16,17}

PATHOPHYSIOLOGY

The effective movement of gastric contents through the stomach depends on 2 major activities: peristalsis of gastric smooth muscle propelling contents to the pylorus and dilation of the pyloric sphincter. Interstitial cells of Cajal (ICCs) are specialized pacemaker cells that drive the contraction of gastric smooth muscle. The speed, strength, and, to a lesser degree, frequency of these contractions are known to be modified by both neurologic and neuroendocrine modulation.

Gastroparesis is defined as the delayed removal of stomach contents in the absence of a physical obstruction.¹⁸ Hyperglycemia mediates nerve damage through a wide range of mechanisms, including poly-adenosine triphosphate ribose, advanced glycosylation end (AGE) products, endoplasmic reticulum stress, oxidative stress, inflammation, and ischemia,^{19,20} which result in demyelination and axonal degeneration.^{21,22} Neuropathy, affecting autonomic input, such as from the vagus nerve, or that affecting the enteric nervous system's own intrinsic neurons, can induce gastroparesis. Loss of parasympathetic stimulation will slow gastric emptying, a picture mimicked during surgical vagotomy.²³ Extensive myelinated and unmyelinated nerve fiber and endoneurial capillary pathologic findings have been found in the vagus nerve of diabetic patients with severe gastroparesis, although interestingly this was comparable to 2 diabetic patients without gastroparesis.²⁴ Both direct neurologic feedback from the small bowel and hormones, such as cholecystikinin and gastric inhibitory peptide, can limit the flow of chyme into the duodenum. Abnormalities of gastric electrical rhythm and transmission can lead to a disruption of the migrating motor complex, ineffective propulsion, and decreased pyloric output.^{25,26} The severity of these arrhythmias has been directly linked to loss of ICCs, although not to the severity of symptoms.²⁷ The origin of gastroparesis has therefore been related to the pathologic features of the ICCs and vagal neurons.^{28,29}

Neural nitric oxide synthase (nNOS) expressed in gastric neurons induces relaxation and accommodation, although its predominant function is to cause dilatation of gastrointestinal sphincters, including the pylorus.²⁹ In animal models of DG, reduced nNOS mRNA, protein, and function³⁰ restricts the passage of food into the duodenum. Interestingly, the reduction in nNOS is not attributable to a loss of active neurons but decreased expression on these intrinsic neurons.⁸

A significant decrease in ICC counts has been found in gastric biopsy specimens of patients with gastroparesis.^{31–34} Interestingly, biopsy specimens of patients with gastroparesis-like syndrome (no evidence of delayed gastric emptying) reveal significantly lower ICC counts,³⁵ emphasizing their critical importance to gastroparesis. Indeed, a decreased ICC count is a key histologic finding in gastroparesis.²⁹ Unlike diabetic neuropathy where the deleterious effects stem from

hyperglycemia, evidence suggests that the decrease in ICC counts is related to the body's response to this condition. Murine models indicate that hyperglycemia per se is not the cause of decreased ICC counts but rather a decrease in the quantity or sensitivity to insulin and insulin-like growth factor.^{36,37}

Murine models using streptozotocin diabetes show a decrease in ICC density and a delay in gastric emptying.³⁸ However, when streptozotocin was given to mice that genetically lacked Csf1, a factor essential for the development of macrophages in gastrointestinal musculature, the effect on ICCs and emptying was lost,³⁹ suggesting a potential role for macrophage-mediated induction of DG. Gastric macrophages differentiate and are classed as M1, a proinflammatory phenotype, or M2, an anti-inflammatory phenotype.⁴⁰ Murine gastroparesis models relatively express larger populations of M1 compared with M2.⁴¹ The presence of the M2 phenotype has been found to protect against gastroparesis in mice through a potent antioxidant, heme oxygenase 1-dependent process.⁴² Up-regulation of heme oxygenase 1 by hemin, a process reproducible in humans, reverses the loss of ICCs, gastric motility, and nNOS in animal models of DG.⁴² In patients with gastroparesis, a loss of CD206⁺ macrophages of the M2 phenotype in the gastric antrum is related to the ICC density.⁴¹ Recent studies show that insulin and insulin-like growth factor 1 can affect macrophage function by promoting inflammation,^{43–45} which may explain how these factors contribute to ICC loss.

Pharmacologic management of diabetes may also affect gastric motility. Endogenous glucagon-like peptide 1 slows gastric emptying⁴⁶; therefore, exogenous glucagon-like peptide 1 analogues and dipeptidyl peptidase 4 inhibitors could theoretically delay gastric emptying. However, clinical studies have not found any effect of these therapies on gastric emptying.^{47–49}

INVESTIGATIONS

The diagnosis of gastroparesis entails the exclusion of a mechanical obstruction, typically with an oesophagogastroduodenoscopy or barium meal. The gold standard for diagnosis is scintigraphy, although breath testing and the SmartPill have been developed as alternatives. Other modalities, such as

ultrasonography, magnetic resonance imaging, and electrogastrography, are much less commonly used.

Gastric-emptying scintigraphy

There has been an attempt to standardize test protocols using gastric-emptying scintigraphy.⁵⁰ A standard low-fat meal is used, although this can be combined with an isotope-labeled liquid. Although a delay in liquid emptying may not be apparent until the development of severe gastroparesis, the sensitivity for detecting gastroparesis may be improved if there is delayed liquid emptying in the presence of normal solid emptying.^{51–53} After an overnight fast, a standard, low-fat, radiolabeled meal is ingested within 10 minutes, and imaging is performed at baseline and 1, 2, and 4 hours with the patient in a standing position. Glucose level should be <275 mg/dL, and use of drugs that affect gastric emptying needs to be discontinued before the procedure. Delayed gastric emptying is defined by >60% retention at 2 hours or >10% retention at 4 hours. A major consideration for this form of imaging is that females can have a physiologic delay in gastric emptying of approximately 15%. Additional pitfalls include intraindividual variation of gastric emptying of up to 24%, and the use of an abnormal low-fat, low-fiber meal that may not mimic real-life meals and thereby lower sensitivity.⁵⁴


Gastric Emptying Breath Test

The gastric emptying breath test relies on the use of a radiolabeled carbon-containing test meal (carbon 13-labeled S platensis or octanoic acid). A similar preparation to gastric-emptying scintigraphy is needed for this investigation. The radiolabeled carbon is released during digestion, and the carbon dioxide is released by respiration through ventilation is measured 4 to 6 hours later. Any exertion that results in increased ventilation rate needs to be avoided.^{54,55} This investigation may be as accurate as gastric-emptying scintigraphy.⁵⁶

Wireless motility capsule

The SmartPill can evaluate pH, pressure, and temperature in addition to whole gut transit time. Data are transmitted directly to a portable receiver. The SmartPill is ingested with a standard meal after use of acid-suppressing medications have been discontinued. Movement of the capsule from an acidic to an alkaline environment represents transit from the stomach to the duodenum, normally within 5 hours. The use of this

Clinical Therapeutics



diagnostic modality allows for ambulatory assessment of patients, is radiation free, and may also provide additional physiologic information.⁵⁴

MANAGEMENT

The general principles of management are to restore nutritional and hydration status, alleviate symptoms, and stabilize diabetes control. The [Figure](#) shows an approach to investigations and management.

Dietary Modifications

In conjunction with improving nutrition, fluid and electrolyte balance also needs to be corrected. Correction of prolonged poor nutrition may predispose patients to refeeding syndrome. Multiple small meals (4 to 6 per day) are preferred to fewer large ones. Meals should be low in fat and fiber because these can slow gastric emptying.^{57–59} Smoking and alcohol consumption may also delay gastric emptying and should be avoided.^{60,61} A trial of 56 patients with DG found that a small particle size diet may alleviate symptoms.⁶² High-calorie liquid drinks may also be a useful adjunct to the management of gastroparesis. Certain foods, such as pizza, orange juice, coffee, broccoli, salsa, and roast beef, which fall into the categories of spicy, acidic, fatty, and high fiber and may delay gastric emptying, should be limited or excluded.⁶³

When oral nutrition is not possible or inadequate, assisted nutrition should be considered. The enteral route is preferable to parenteral nutrition when possible because of the lower risk of complications,

such as line infection and thrombosis. A nasojejun tube may be used initially, followed by a jejunostomy tube, if required.⁵⁹

Glycemic control

The association between glycemic control and gastroparesis is not fully understood and, of course, may be bidirectional. Earlier studies found that acute hyperglycemia delayed gastric emptying in healthy individuals.^{64,65} Subsequent studies found that in people with T1DM, hyperglycemia prolonged gastric emptying of solids and liquids,^{66–68} with reduced antral mobility and increased proximal gastric adherence.^{67,69} In contrast, acute hypoglycemia accelerates gastric emptying in healthy controls.⁷⁰ Paradoxically, in patients with T2DM, a higher fasting blood glucose level was associated with faster gastric emptying.^{71,72}

There are conflicting data on the effect of long-term glycemic control on gastric emptying, with several studies having previously found no correlation in patients with T2DM and T2DM.^{71,73–75} However, recently, in a follow-up cohort of 78 patients from the Diabetes Control and Complications Trial/Epidemiology of Diabetes Interventions and Complications studies who underwent an assessment of gastric emptying using the breath test, baseline HbA_{1c} level, duration of diabetes, and mean HbA_{1c} level during the trial period were associated with delayed gastric emptying.⁷⁶ A further retrospective study of patients with diabetes who had undergone gastric-emptying scintigraphy found that a higher HbA_{1c} level was significantly associated with higher gastric retention at 4 hours.⁷⁷

PHARMACOLOGIC TREATMENT

Prokinetics, used to promote gastrointestinal tract motility, have been and remain the mainstay of treatment for gastroparesis ([Table](#)).⁷⁸ However, data are lacking on the long-term effectiveness of these drugs, perhaps also reflecting the varying clinical course of this condition whereby long-term treatment is often not necessary.⁷

Metoclopramide

Metoclopramide is a potent dopamine D₂ receptor antagonist and serotonin (5HT₄) receptor agonist that acts on the brainstem and peripheral nerves. In the

Table. Pharmacological treatment options for the management of diabetic gastroparesis.

Class of Medication	Effect on GI Tract	Use in Diabetic Gastroparesis
Dopamine D ₂ receptor antagonists (metoclopramide, domperidone) approved for diabetic gastroparesis. Domperidone may avoid CNS adverse effects.	Increases contractions of the gastric antrum by releasing acetylcholine from enteric neurons	Metoclopramide is FDA
Motilin receptor agonists (erythromycin, mitemincal, and camicinal)	Increased antral contraction	Erythromycin favored in extremely unwell, hospitalized patients, risk of tachyphylaxis with erythromycin
5HT ₄ receptor agonists (cisapride, tegaserod, revexepride)	Increased muscular contraction via cholinergic pathways	Older agents not used due to cardiac safety concerns, more selective agents undergoing evaluation
Ghrelin receptor agonists (TZP 101, TZP 102, relamorelin)	Increased migrating motor complexes (Phase III) and vagal signaling.	Significant improvement in symptoms in clinical trials
Neurokinin receptor antagonists (aprepitant)	Antiemetic, antagonizes the effects of substance P	Limited trial data

CNS = central nervous system; FDA = US Food and Drug Administration; GI = gastrointestinal.

gastrointestinal tract, it increases contractions of the gastric antrum by releasing acetylcholine from enteric neurons.^{79–81} The central effects may account in some part for the reduction of nausea. An early double-blind, placebo-controlled trial of 40 patients with DG found that metoclopramide 10 mg QID PO produced a significant improvement in meal tolerance, symptoms of gastroparesis, and gastric emptying during 3 weeks.⁸² In a subsequent double-blind, placebo-controlled study of 13 patients with DG, metoclopramide was administered parenterally and then orally before each meal and before bed and produced a mean symptom reduction of 52.6% and improved gastric emptying in 7 patients.⁸³ It remains the only drug approved by the US Food and Drug Administration (FDA) for the treatment of gastroparesis. However, concerns regarding potentially irreversible tardive dyskinesia has led to an FDA warning restricting its use to no longer than 3 months and recommendations to use the lowest effective dose for the shortest possible time. Indeed, along with older age and female sex, diabetes itself is a risk factor for developing tardive dyskinesia.⁸¹

Domperidone

Domperidone, another dopamine D₂ receptor antagonist, does not cross the blood brain barrier and therefore does not cause the same central nervous system adverse effects.⁸⁴ A dose of 10 mg TID improves a range of gastroparesis-related symptoms.⁸⁵ This drug is not currently FDA approved, and its prescription in the United States requires an investigational new drug application because of the risk of QTc prolongation and cardiac arrhythmias.

Motilin Receptor Agonists

Erythromycin is a macrolide antibiotic that agonizes motilin receptors, leading to increased antral contraction.⁸⁶ The intravenous route is strongly favored in hospitalized patients, although tachyphylaxis may occur with this drug usually after 4 weeks of use.⁸⁷ There are potential interactions with other drugs because of cytochrome P450 C3A4 inhibition. Clarithromycin and azithromycin have also been used, but there is a paucity of supporting clinical trial data. A retrospective case-control analysis

of 120 patients found equal efficacy between azithromycin and erythromycin in accelerating gastric emptying in gastroparesis.⁸⁸ A smaller study that evaluated intravenous use of erythromycin and azithromycin found similar stimulation of antral activity with a longer duration of effect.⁸⁹ The advantages of azithromycin are the lack of P450 inhibition, longer duration of action, and better adverse effect profile; however, more research needs to be performed to investigate its utility further. Macrolides can also cause QTc prolongation through their effect on I_{Kr} potassium channels.⁹⁰

New agents acting on the motilin receptor, including mitemincal and camicinal, deliver the benefits of treatment without the antibiotic activity reducing the risk of tachyphylaxis. However, a double-blind, placebo-controlled trial of mitemincal found no symptom relief compared with placebo despite an improvement in gastric emptying.⁹¹ A single dose of camicinal in patients with T1DM significantly accelerates gastric emptying of solids, and further trials are under way.⁹²

5HT₄ Receptor Agonists

The 5HT₄ receptor agonist class of drugs activate 5HT₄ receptors, leading to increased muscular contraction via cholinergic pathways.⁹³ Cisapride, a 5HT₄ receptor agonist and established drug for gastroparesis, was withdrawn because of cardiac safety concerns attributable to activation of hERG potassium channels, leading to QTc prolongation and ventricular arrhythmias. Similarly, tegaserod, another 5HT₄ agonist, has been withdrawn because of cardiac safety concerns.⁹⁴ Recently, more selective 5HT₄ receptor agonists have been developed, but a trial of revexepride in patients with gastroparesis found no benefit on symptoms or gastric emptying versus placebo.⁹⁵

Ghrelin Receptor Agonists

Ghrelin is a peptide released from gastric mucosal endocrine cells.⁹⁶ Stimulation of the GHS-R1a receptor results in increased migrating motor complexes (Phase III) and vagal signaling.^{97,98} Four single daily intravenous infusions of TZP 101 (ulimorelin) were associated with a significant improvement in symptoms of gastroparesis up to 30 days after administration in 23 patients with DG compared with placebo.⁹⁹ In a Phase IIa trial of 92 patients with DG, TZP 102, an oral ghrelin receptor agonist, produced a

significant improvement in symptoms during 28 days.¹⁰⁰ A subsequent Phase IIb study of TZP 102 in 201 patients with DG found significant symptom improvement but failed to establish efficacy compared with placebo.¹⁰¹ Recently, relamorelin, a subcutaneously administered ghrelin agonist, also produced a significant improvement in vomiting and gastric emptying compared with placebo in patients with DG.¹⁰²

Neurokinin Receptor Antagonists

Substance P is a peptide involved in the induction of vomiting, with actions through binding to neurokinin 1 receptors.¹⁰³ Aprepitant is a neurokinin 1 receptor antagonist that is currently widely used in chemotherapy-induced vomiting and nausea. Fountoulakis reported 2 cases of DG that were successfully treated with aprepitant for 12 and 18 months.¹⁰⁴ There are currently no published trials of this drug for DG.

OTHER THERAPIES

Botulinum Toxin

A manifestation of DG is pylorospasm, and botulinum toxin, which blocks acetylcholine release at the neuromuscular junction and reduces excessive pyloric contraction, has been evaluated.^{105–107} Initial case series showed an improvement in symptoms and gastric emptying^{108–110}; however, 2 small randomized controlled trials have not yielded positive results,^{111,112} questioning the efficacy of this treatment in gastroparesis.¹¹³ Furthermore, a retrospective analysis of 179 cases treated with botulinum toxin suggested that diabetes further limited a benefit.¹¹⁴

Transpyloric Stenting

Endoscopic stent placement has been investigated in small series of patients with gastroparesis. Khashab et al¹¹⁵ reported their experience in 30 patients undergoing a total of 48 procedures. Although the data were incomplete, there were reports of symptom improvement and more rapid gastric emptying, but stent migration is a complication. The authors suggested that transpyloric stenting may be a useful salvage therapy or may help identify patients who may benefit from other therapies directed at the pylorus.

Gastric Electrical Stimulation

Gastric electrical stimulation treatment has been approved by the FDA and the National Institute for Health and Care Excellence for the treatment of drug refractory DG. A device implanted into the abdominal wall has leads that extend into the greater curvature of the stomach and generates 12 pulses per minute. Abell et al¹¹⁶ studied 38 patients undergoing gastric electrical stimulation and found a reduction in symptoms with associated weight gain. Long-term follow-up has reported morbidity reduction for up to 10 years after the procedure.^{116,117} Heckert et al¹¹⁸ reported an improvement in symptoms in 75% of 151 patients undergoing this procedure, and patients with diabetes surprisingly had greater success.

Gastric Per-Oral Endoscopic Myotomy

Khashab et al¹¹⁹ reported an improvement in symptoms in 26 of 30 patients undergoing gastric per-oral endoscopic myotomy for refractory gastroparesis. However, in patients undergoing this procedure for mechanical obstruction, there was a high rate of dumping syndrome with postprandial hypoglycemia, but this appears less common in patients with gastroparesis.¹²⁰

SURGICAL TREATMENTS

A number of treatments have been used in severe intractable gastroparesis. Laparoscopic pyloroplasty was evaluated retrospectively in 46 patients and revealed a significant symptom improvement and gastric emptying in 90% of cases.¹²¹ Gastrectomy has also been used in selected patients to alleviate severe refractory symptoms, may be considered in those with high risk of renal failure or premature death,^{122–124} and may be considered earlier in patients with failed GES.¹²⁵ Pancreatic transplantation may be of benefit in DG, perhaps in the short term via improved glycemic control and in the long term through nerve fiber regeneration.^{126,127}

APPROACH TO THE MANAGEMENT OF GASTROPARESIS

The American College of Gastroenterology guidelines recommend that a documented delay in gastric emptying is required for the diagnosis, with scintigraphy being the most reliable gold standard.¹¹³ Other alternatives still require further validation for use in

the diagnosis of DG. Before undertaking any investigation, use of all medications that have the potential to affect gastric emptying must be stopped for at least 48 hours. These medications include narcotic opioid analgesics and anticholinergic agents (may give a falsely delayed result) or the medications metoclopramide, domperidone, and erythromycin (which accelerate gastric emptying and may give a falsely normal result). Hyperglycemia can lead to a deterioration in symptoms, so glycemic control should be measured and improved before testing during any gastric emptying studies.

Primarily, the management of DG is ensuring hydration, adequate electrolytes, and nutritional support. One of the main themes of any guidelines on gastroparesis is the importance of optimizing glycemic control. Dietician input should be sought with regard to the consumption of frequent small-volume meals low in fat and soluble fiber. The guidelines suggest that indications for enteral nutrition include unintentional loss of $\geq 10\%$ of usual weight during 3 to 6 months and/or repeated admission for refractory symptoms. A nasogastric tube should be used initially, followed by a jejunostomy if indicated. This approach may reduce hospital admissions and improve symptoms.¹²⁸

The strongest evidence and recommendations are with metoclopramide as the first line of prokinetic therapy, which should be given at the lowest effective dose in liquid formation to facilitate absorption. It is approved for a duration of 12 weeks, and advice should be given to discontinue therapy if patients develop adverse effects, such as tardive dyskinesia. If patients do not tolerate metoclopramide, then domperidone can be used with investigational new drug clearance from the FDA. Domperidone does not have the predisposition for central nervous system adverse effects as metoclopramide does; however, there is a tendency for this to cause a prolonged QTc in interval, so a baseline ECG is recommended. If the QTc is > 470 msec in men and 450 msec in women, then this should be withheld. A follow-up ECG while the patient is receiving domperidone treatment is advised. The use of erythromycin orally and intravenously in hospital when intravenous prokinetic therapy is initiated is recommended.

The surgical procedures mentioned earlier may be considered on an individual basis in refractory cases. In patients in whom nausea and vomiting are an issue,

antiemetics should be considered for symptomatic treatment, and in refractory cases, tricyclic antidepressants can be used; however, caution must be taken they may retard gastric emptying. The National Institute for Health and Care Excellence guidelines advise that gastric electrical stimulation is an option patients with chronic, intractable nausea and vomiting, and its use in particular is beneficial in DG.¹²⁹

CONCLUSION

DG remains a challenging complication of diabetes. Advances in understanding the pathophysiology of this condition have been limited. The management of gastroparesis is difficult because there is discordance between symptoms and the degree of delayed gastric emptying. Because of the relatively low prevalence of DG and the fluctuating symptoms, it is often difficult to recruit patients into clinical trials. Current treatment options are limited; however, recent pharmacologic and endoscopic interventions show promise.

CONFLICTS OF INTEREST

The authors have indicated that they have no conflicts of interest regarding the content of this article.

REFERENCES

1. Rundles R. Diabetic Neuropathy. *Medicine (Baltimore)*. 1945;24:111–160.
2. Kassander P. Asymptomatic gastric retention in diabetics (gastroparesis diabetorum). *Ann Intern Med*. 1958;48:797–812.
3. Bharucha AE. Epidemiology and natural history of gastroparesis. *Gastroenterol Clin North Am*. 2015;44:9–19.
4. Camilleri M. Novel Diet, Drugs, and Gastric Interventions for Gastroparesis. *Clin Gastroenterol Hepatol*. 2016;14:1072–1080.
5. Teigland T, et al. A longitudinal study on patients with diabetes and symptoms of gastroparesis - associations with impaired quality of life and increased depressive and anxiety symptoms. *J Diabetes Complications*. 2018;32:89–94.
6. Homko C, Siraj ES, Parkman HP. The impact of gastroparesis on diabetes control: Patient perceptions. *J Diabetes Complications*. 2016;30:826–829.
7. Alam U, Asghar O, Malik RA. Diabetic gastroparesis: Therapeutic options. *Diabetes Ther*. 2010;1:32–43.
8. Camilleri M, Bharucha AE, Farrugia G. Epidemiology, mechanisms, and management of diabetic gastroparesis. *Clin Gastroenterol Hepatol*. 2011;9:5–12. quiz e7.
9. Lysy J, et al. Relationships between hypoglycaemia and gastric emptying abnormalities in insulin-treated diabetic patients. *Neurogastroenterol Motil*. 2006;18:433–440.
10. Horowitz M, et al. 'Gastric' hypoglycaemia—an important concept in diabetes management. *Neurogastroenterol Motil*. 2006;18:405–407.
11. Choung RS, et al. Risk of gastroparesis in subjects with type 1 and 2 diabetes in the general population. *Am J Gastroenterol*. 2012;107:82–88.
12. Aleppo G, et al. Reported gastroparesis in adults with type 1 diabetes (T1D) from the T1D Exchange clinic registry. *J Diabetes Complications*. 2017;31:1669–1673.
13. Maleki D, et al. Gastrointestinal tract symptoms among persons with diabetes mellitus in the community. *Arch Intern Med*. 2000;160:2808–2816.
14. Bytzer P, et al. Prevalence of gastrointestinal symptoms associated with diabetes mellitus: a population-based survey of 15,000 adults. *Arch Intern Med*. 2001;161:1989–1996.
15. Bharucha AE, et al. Relationship between clinical features and gastric emptying disturbances in diabetes mellitus. *Clin Endocrinol (Oxf)*. 2009;70:415–420.
16. Talley NJ, et al. Predictors of turnover of lower gastrointestinal symptoms in diabetes mellitus. *Am J Gastroenterol*. 2002;97:3087–3094.
17. Quan C, et al. Gain and loss of gastrointestinal symptoms in diabetes mellitus: associations with psychiatric disease, glycemic control, and autonomic neuropathy over 2 years of follow-up. *Am J Gastroenterol*. 2008;103:2023–2030.
18. Parkman HP, et al. American Gastroenterological Association technical review on the diagnosis and treatment of gastroparesis. *Gastroenterology*. 2004;127:1592–1622.
19. Albers JW, Pop-Busui R. Diabetic neuropathy: mechanisms, emerging treatments, and subtypes. *Curr Neurol Neurosci Rep*. 2014;14:473.
20. Hussain N, Adrian TE. Diabetic Neuropathy: Update on Pathophysiological Mechanism and the Possible Involvement of Glutamate Pathways. *Curr Diabetes Rev*. 2017;13:488–497.
21. Guy RJ, et al. Diabetic gastroparesis from autonomic neuropathy: surgical considerations and changes in vagus nerve morphology. *J Neurol Neurosurg Psychiatry*. 1984;47:686–691.
22. Smith B. Neuropathology of the oesophagus in diabetes mellitus. *J Neurol Neurosurg Psychiatry*. 1974;37:1151–1154.
23. Shafi MA, Pasricha PJ. Post-surgical and obstructive gastroparesis. *Curr Gastroenterol Rep*. 2007;9:280–285.
24. Britland ST, et al. Vagus nerve morphology in diabetic gastropathy. *Diabet Med*. 1990;7:780–787.
25. O'Grady G, Abell T. Gastric arrhythmias in gastroparesis: low and high resolution mapping of gastric electrical activity. *Gastroenterol Clin North Am*. 2015;44:169–184.

26. Waseem S, Moshiree B, Dragano PV. Gastroparesis: Current diagnostic challenges and management considerations. *World J Gastroenterol*. 2009;15:25–37.
27. Lin Z, et al. Association of the status of interstitial cells of Cajal and electrogastrogram parameters, gastric emptying and symptoms in patients with gastroparesis. *Neurogastroenterol Motil*. 2010;22:56–61. e10.
28. Camilleri M, Bharucha AE, Farugia G. Epidemiology, Mechanisms and Management of Diabetic Gastroparesis. *Clin Gastroenterol Hepatol*. 2011;9:5–e7.
29. Jung Hwan O. Recent Advances in the Pathophysiology and Treatment of Gastroparesis. *J Neurogastroenterol Motil*. 2013;19:18–24.
30. Watkins CC, et al. Insulin restores neuronal nitric oxide synthase expression and function that is lost in diabetic gastropathy. *J Clin Invest*. 2000;106:803.
31. Harberson J, et al. Gastric neuromuscular pathology in gastroparesis: analysis of full-thickness antral biopsies. *Dig Dis Sci*. 2010;55:359–370.
32. Iwasaki H, et al. A deficiency of gastric interstitial cells of Cajal accompanied by decreased expression of neuronal nitric oxide synthase and substance P in patients with type 2 diabetes mellitus. *J Gastroenterol*. 2006;41:1076–1087.
33. McCallum RW, et al. Su1455 Relationship of Gastric Emptying With Numbers of Interstitial Cells of Cajal in Antral Muscularis Propria of Gastroparesis Patient. *Gastroenterology*. 2015;148:S–517.
34. Bernard CE, et al. Association of low numbers of CD206-positive cells with loss of ICC in the gastric body of patients with diabetic gastroparesis. *Neurogastroenterol Motil*. 2014;26:1275–1284.
35. Bashashati M, et al. Pathological Findings of the Antral and Pyloric Smooth Muscle in Patients with Gastroparesis-Like Syndrome Compared to Gastroparesis: Similarities and Differences. *Dig Dis Sci*. 2017;62:2828–2833.
36. Horvath VJ, Vittal H, Ordog T. Reduced insulin and IGF-I signaling, not hyperglycemia, underlies the diabetes-associated depletion of interstitial cells of Cajal in the murine stomach. *Diabetes*. 2005;54:1528–1533.
37. Yang S, et al. Impaired insulin/IGF-1 is responsible for diabetic gastroparesis by damaging myenteric cholinergic neurones and interstitial cells of Cajal. *Biosci Rep*. 2017;37:5.
38. Wang XY, et al. Loss of intramuscular and submuscular interstitial cells of Cajal and associated enteric nerves is related to decreased gastric emptying in streptozotocin-induced diabetes. *Neurogastroenterol Motil*. 2009;21. 1095–e92.
39. Cipriani G, et al. Diabetic Csf1 (op/op) mice lacking macrophages are protected against the development of delayed gastric emptying. *Cell Mol Gastroenterol Hepatol*. 2016;2:40–47.
40. Srinivasan S. Macrophages: The Missing Link in Diabetic Gastroparesis? *Cell Mol Gastroenterol Hepatol*. 2016;2:5–6.
41. Choi KM, et al. CD206-positive M2 macrophages that express heme oxygenase-1 protect against diabetic gastroparesis in mice. *Gastroenterology*. 2010;138:2399–2409. 2409.e1.
42. Choi KM, et al. Heme oxygenase-1 protects interstitial cells of Cajal from oxidative stress and reverses diabetic gastroparesis. *Gastroenterology*. 2008;135:2055–2064. 2064.e1–2.
43. Han CZ, et al. Macrophages redirect phagocytosis by non-professional phagocytes and influence inflammation. *Nature*. 2016;539:570.
44. Manowsky J, et al. Insulin-induced cytokine production in macrophages causes insulin resistance in hepatocytes. *Am J Physiol Endocrinol Metab*. 2016;310:E938–E946.
45. Spadaro O. IGF1 shapes the macrophage activation in response to immunometabolic challenge. 2017;19:225–234.
46. Marathe CS, et al. Relationships Between Gastric Emptying, Postprandial Glycemia, and Incretin Hormones. *Diabetes Care*. 2013;36:1396–1405.
47. DeFronzo RA, et al. Effects of exenatide versus sitagliptin on postprandial glucose, insulin and glucagon secretion, gastric emptying, and caloric intake: a randomized, cross-over study. *Curr Med Res Opin*. 2008;24:2943–2952.
48. Stevens JE, et al. The effects of sitagliptin on gastric emptying in healthy humans - a randomised, controlled study. *Aliment Pharmacol Ther*. 2012;36:379–390.
49. Delgado-Aros S, et al. Effects of glucagon-like peptide-1 and feeding on gastric volumes in diabetes mellitus with cardiovascular dysfunction. *Neurogastroenterol Motil*. 2003;15:435–443.
50. Abell TL, et al. Consensus recommendations for gastric emptying scintigraphy: a joint report of the American Neurogastroenterology and Motility Society and the Society of Nuclear Medicine. *Am J Gastroenterol*. 2008;103:753–763.
51. Loo FD, et al. Gastric emptying in patients with diabetes mellitus. *Gastroenterology*. 1984;86:485–494.
52. Ziessman HA, et al. The added diagnostic value of liquid gastric emptying compared with solid emptying alone. *J Nucl Med*. 2009;50:726–731.

53. Sachdeva P, et al. Gastric emptying of solids and liquids for evaluation for gastroparesis. *Dig Dis Sci*. 2011;56:1138–1146.
54. Shin AS, Camilleri M. Diagnostic assessment of diabetic gastroparesis. *Diabetes*. 2013;62:2667–2673.
55. Navas CM, Patel NK, Lacy BE. Gastroparesis: Medical and Therapeutic Advances. *Dig Dis Sci*. 2017;62:2231–2240.
56. Bharucha AE, et al. Comprehensive assessment of gastric emptying with a stable isotope breath test. *Neurogastroenterol Motil*. 2013;25:e60–e69.
57. Bouras EP, Scolapio JS. Gastric motility disorders: management that optimizes nutritional status. *J Clin Gastroenterol*. 2004;38:549–557.
58. Parrish CR YC. Nutritional intervention for the patient with gastroparesis: an update. *Pract Gastroenterol*. 2005;29:29–66.
59. Liu N, Abell T. Gastroparesis Updates on Pathogenesis and Management. *Gut Liver*. 2017;11:579–589.
60. Sanaka M, et al. Effect of cigarette smoking on gastric emptying of solids in Japanese smokers: a crossover study using the ¹³C-octanoic acid breath test. *J Gastroenterol*. 2005;40:578–582.
61. Pfeiffer A, Hög B, Kaess H. Effect of ethanol and commonly ingested alcoholic beverages on gastric emptying and gastrointestinal transit. *Clin Invest*. 1992;70:487–491.
62. Olausson EA, et al. A small particle size diet reduces upper gastrointestinal symptoms in patients with diabetic gastroparesis: a randomized controlled trial. *Am J Gastroenterol*. 2014;109:375–385.
63. Wytiacz V, et al. Foods provoking and alleviating symptoms in gastroparesis: patient experiences. *Dig Dis Sci*. 2015;60:1052–1058.
64. Stunkard AJ, Wolff HG. Studies on the physiology of hunger. I. The effect of intravenous administration of glucose on gastric hunger contractions in man. *J Clin Invest*. 1956;35:954–963.
65. MacGregor IL, et al. The effect of acute hyperglycemia on gastric emptying in man. *Gastroenterology*. 1976;70:190–196.
66. Fraser RJ, et al. Hyperglycaemia slows gastric emptying in Type 1 (insulin-dependent) diabetes mellitus. *Diabetologia*. 1990;33:675–680.
67. Samsom M, et al. Gastrointestinal motor mechanisms in hyperglycaemia induced delayed gastric emptying in type I diabetes mellitus. *Gut*. 1997;40:641–646.
68. Schvarcz E, et al. Physiological hyperglycemia slows gastric emptying in normal subjects and patients with insulin-dependent diabetes mellitus. *Gastroenterology*. 1997;113:60–66.
69. Rayner CK, et al. Proximal gastric compliance and perception of distension in type 1 diabetes mellitus: effects of hyperglycemia. *Am J Gastroenterol*. 2000;95:1175–1183.
70. Schvarcz E, et al. Atropine inhibits the increase in gastric emptying during hypoglycemia in humans. *Diabetes Care*. 1995;18:1463–1467.
71. Bharucha AE, et al. Relationship between glycemic control and gastric emptying in poorly controlled type 2 diabetes. *Clin Gastroenterol Hepatol*. 2015;13:466–476. e1.
72. Halland M, Bharucha AE. Relationship Between Control of Glycemia and Gastric Emptying Disturbances in Diabetes Mellitus. *Clin Gastroenterol Hepatol*. 2016;14:929–936.
73. Hyett B, et al. Delayed radionuclide gastric emptying studies predict morbidity in diabetics with symptoms of gastroparesis. *Gastroenterology*. 2009;137:445–452.
74. Merio R, et al. Slow gastric emptying in type I diabetes: relation to autonomic and peripheral neuropathy, blood glucose, and glycemic control. *Diabetes Care*. 1997;20:419–423.
75. Reddy S, et al. Do HbA1C Levels Correlate With Delayed Gastric Emptying in Diabetic Patients? *J Neurogastroenterol Motil*. 2010;16:414–417.
76. Bharucha AE, et al. Delayed Gastric Emptying Is Associated With Early and Long-term Hyperglycemia in Type 1 Diabetes Mellitus. *Gastroenterology*. 2015;149:330–339.
77. Izzy M, et al. Glycosylated hemoglobin level may predict the severity of gastroparesis in diabetic patients. *Diabetes Res Clin Pract*. 2017;135:45–49.
78. Perkel MS, et al. Metoclopramide therapy in patients with delayed gastric emptying: a randomized, double-blind study. *Dig Dis Sci*. 1979;24:662–666.
79. Peringer E, Jenner P, Marsden CD. Effect of metoclopramide on turnover of brain dopamine noradrenaline and 5-hydroxytryptamine. *J Pharm Pharmacol*. 1975;27:442–444.
80. Koch KL. Diabetic gastropathy: gastric neuromuscular dysfunction in diabetes mellitus: a review of symptoms, pathophysiology, and treatment. *Dig Dis Sci*. 1999;44:1061–1075.
81. Pasricha PJ, et al. Drug Insight: from disturbed motility to disordered movement—a review of the clinical benefits and medicolegal risks of metoclopramide. *Nat Clin Pract Gastroenterol Hepatol*. 2006;3:138–148.

82. McCallum RW, et al. A multicenter placebo-controlled clinical trial of oral metoclopramide in diabetic gastroparesis. *Diabetes Care*. 1983;6:463-467.
83. Ricci DA, et al. Effect of metoclopramide in diabetic gastroparesis. *J Clin Gastroenterol*. 1985;7:25-32.
84. Patterson D, et al. A double-blind multicenter comparison of domperidone and metoclopramide in the treatment of diabetic patients with symptoms of gastroparesis. *Am J Gastroenterol*. 1999;94:1230-1234.
85. Heckert J, Parkman HP. Therapeutic response to domperidone in gastroparesis: A prospective study using the GCSI-daily diary. *Neurogastroenterol Motil*. 2018;30:1.
86. Janssens J, et al. Improvement of gastric emptying in diabetic gastroparesis by erythromycin. Preliminary studies. *N Engl J Med*. 1990;322:1028-1031.
87. Richards RD, Davenport K, McCallum RW. The treatment of idiopathic and diabetic gastroparesis with acute intravenous and chronic oral erythromycin. *Am J Gastroenterol*. 1993;88:203-207.
88. Larson JM, et al. Advantages of azithromycin over erythromycin in improving the gastric emptying half-time in adult patients with gastroparesis. *J Neurogastroenterol Motil*. 2010;16:407-413.
89. Moshiree B, et al. Comparison of the effect of azithromycin versus erythromycin on antroduodenal pressure profiles of patients with chronic functional gastrointestinal pain and gastroparesis. *Dig Dis Sci*. 2010;55:675-683.
90. Albert RK, Schuller JL. Macrolide antibiotics and the risk of cardiac arrhythmias. *Am J Respir Crit Care Med*. 2014;189:1173-1180.
91. McCallum RW, Cynshi O. Clinical trial: effect of mitemincin (a motilin agonist) on gastric emptying in patients with gastroparesis - a randomized, multicentre, placebo-controlled study. *Aliment Pharmacol Ther*. 2007;26:1121-1130.
92. Hellstrom PM, et al. The pharmacodynamics, safety and pharmacokinetics of single doses of the motilin agonist, camicinal, in type 1 diabetes mellitus with slow gastric emptying. *Br J Pharmacol*. 2016;173:1768-1777.
93. McCallum RW, et al. Cisapride. A preliminary review of its pharmacodynamic and pharmacokinetic properties, and therapeutic use as a prokinetic agent in gastrointestinal motility disorders. *Drugs*. 1988;36:652-681.
94. Quigley EMM. Prokinetics in the Management of Functional Gastrointestinal Disorders. *Curr Gastroenterol Rep*. 2017;19:53.
95. Tack J, et al. Randomized clinical trial: a controlled pilot trial of the 5-HT₄ receptor agonist revexepide in patients with symptoms suggestive of gastroparesis. *Neurogastroenterol Motil*. 2016;28:487-497.
96. Date Y, et al. Ghrelin, a novel growth hormone-releasing acylated peptide, is synthesized in a distinct endocrine cell type in the gastrointestinal tracts of rats and humans. *Endocrinology*. 2000;141:4255-4261.
97. Tack J, et al. Influence of ghrelin on interdigestive gastrointestinal motility in humans. *Gut*. 2006;55:327-333.
98. Page AJ, et al. Ghrelin selectively reduces mechanosensitivity of upper gastrointestinal vagal afferents. *Am J Physiol Gastrointest Liver Physiol*. 2007;292:G1376-G1384.
99. Wo JM, et al. Randomised clinical trial: ghrelin agonist TZP-101 relieves gastroparesis associated with severe nausea and vomiting-randomised clinical study subset data. *Aliment Pharmacol Ther*. 2011;33:679-688.
100. Ejskjaer N, et al. A phase 2a, randomized, double-blind 28-day study of TZP-102 a ghrelin receptor agonist for diabetic gastroparesis. *Neurogastroenterol Motil*. 2013;25:e140-e150.
101. McCallum RW, et al. Phase 2b, randomized, double-blind 12-week studies of TZP-102, a ghrelin receptor agonist for diabetic gastroparesis. *Neurogastroenterol Motil*. 2013;25:e705-e717.
102. Lembo A, et al. Relamorelin Reduces Vomiting Frequency and Severity and Accelerates Gastric Emptying in Adults With Diabetic Gastroparesis. *Gastroenterology*. 2016;151:87-96. e6.
103. Mantyh PW. Neurobiology of substance P and the NK1 receptor. *J Clin Psychiatry*. 2002;63:6-10.
104. Fountoulakis N, et al. Successful management of refractory diabetic gastroparesis with long-term Aprepitant treatment. *Diabet Med*. 2017;34:1483-1486.
105. Mearin F, Camilleri M, Malagelada JR. Pyloric dysfunction in diabetics with recurrent nausea and vomiting. *Gastroenterology*. 1986;90:1919-1925.
106. Camilleri M, Brown ML, Malagelada JR. Relationship between impaired gastric emptying and abnormal gastrointestinal motility. *Gastroenterology*. 1986;91:94-99.
107. Jankovic J, Brin MF. Therapeutic uses of botulinum toxin. *N Engl J Med*. 1991;324:1186-1194.
108. Ezzeddine D, et al. Pyloric injection of botulinum toxin for treatment of diabetic gastroparesis. *Gastrointest Endosc*. 2002;55:920-923.
109. Lacy BE, et al. Botulinum toxin for the treatment of gastroparesis: a preliminary report. *Am J Gastroenterol*. 2002;97:1548-1552.
110. Bromer MQ, et al. Endoscopic pyloric injection of botulinum toxin A for the treatment of refractory gastroparesis. *Gastrointest Endosc*. 2005;61:833-839.

111. Arts J, et al. Clinical trial: a randomized-controlled crossover study of intrapyloric injection of botulinum toxin in gastroparesis. *Aliment Pharmacol Ther.* 2007;26:1251–1258.
112. Friedenberg FK, et al. Botulinum toxin A for the treatment of delayed gastric emptying. *Am J Gastroenterol.* 2008;103:416–423.
113. Camilleri M PH, Shafi MA, Abell TA, Gerson L. Clinical Guideline: Management of Gastroparesis. *Am J Gastroenterol.* 2013;108:18–38.
114. Coleski R, Anderson MA, Hasler WL. Factors associated with symptom response to pyloric injection of botulinum toxin in a large series of gastroparesis patients. *Dig Dis Sci.* 2009;54:2634–2642.
115. Khashab MA, et al. Refractory gastroparesis can be successfully managed with endoscopic transpyloric stent placement and fixation (with video). *Gastrointest Endosc.* 2015;82:1106–1109.
116. Abell TL, et al. Gastric electrical stimulation in intractable symptomatic gastroparesis. *Digestion.* 2002;66:204–212.
117. McCallum RW, et al. Gastric electrical stimulation improves outcomes of patients with gastroparesis for up to 10 years. *Clin Gastroenterol Hepatol.* 2011;9:314–319. e1.
118. Heckert J, et al. Gastric Electric Stimulation for Refractory Gastroparesis: A Prospective Analysis of 151 Patients at a Single Center. *Dig Dis Sci.* 2016;61:168–175.
119. Khashab MA, et al. Gastric peroral endoscopic myotomy for refractory gastroparesis: results from the first multicenter study on endoscopic pyloromyotomy (with video). *Gastrointest Endosc.* 2017;85:123–128.
120. Louazon T, et al. Temporary dumping syndrome after gastric peroral endoscopic myotomy: should we control the glycemia? *Endoscopy.* 2016;48:E10–E11.
121. Mancini SA, et al. Pyloroplasty for Refractory Gastroparesis. *Am Surg.* 2015;81:738–746.
122. Ejskjaer NT, et al. Novel surgical treatment and gastric pathology in diabetic gastroparesis. *Diabet Med.* 1999;16:488–495.
123. Watkins PJ, Buxton-Thomas MS, Howard ER. Long-term outcome after gastrectomy for intractable diabetic gastroparesis. *Diabet Med.* 2003;20:58–63.
124. Bhayani NH, et al. End of the road for a dysfunctional end organ: laparoscopic gastrectomy for refractory gastroparesis. *J Gastrointest Surg.* 2015;19:411–417.
125. Zehetner J, et al. Minimally invasive surgical approach for the treatment of gastroparesis. *Surg Endosc.* 2013;27:61–66.
126. Cashion AK, et al. Gastroparesis following kidney/pancreas transplant. *Clin Transplant.* 2004;18:306–311.
127. Mehra S, et al. Corneal confocal microscopy detects early nerve regeneration after pancreas transplantation in patients with type 1 diabetes. *Diabetes Care.* 2007;30:2608–2612.
128. Fontana RJ, Barnett JL. Jejunostomy tube placement in refractory diabetic gastroparesis: a retrospective review. *Am J Gastroenterol.* 1996;91:2174–2178.
129. O'Grady G, et al. High-frequency gastric electrical stimulation for the treatment of gastroparesis: a meta-analysis. *World J Surg.* 2009;33:1693–1701.

Address correspondence to: Shazli Azmi, PhD, Manchester University Hospital NHS Foundation Trust, Manchester Diabetes Centre, Manchester Royal Infirmary, 193 Hathersage Road, Manchester M13 0JE. E-mail: Shazli.azmi@manchester.ac.uk